

A Recent Look at the Quantum Berezinian in the Yangian $Y(\mathfrak{gl}_{m|n})$

Lucy Gow*
University of Sydney

February 1, 2008

Brundan and Kleshchev recently introduced a new family of presentations of the Yangian $Y(\mathfrak{gl}_n)$ associated to the general linear Lie algebra $\mathfrak{gl}_n$, and thus provided a fresh approach to its study. In this article, we would like to show how some of their ideas can be fruitfully extended to consider the Yangian $Y(\mathfrak{gl}_{m|n})$ associated to the Lie superalgebra $\mathfrak{gl}_{m|n}$. In particular, we give a new proof of the result by Nazarov that the quantum Berezinian is central.¹

1 Definition of $Y(\mathfrak{gl}_{m|n})$

The Yangian $Y(\mathfrak{gl}_{m|n})$ is defined in [7] to be the $\mathbb{Z}_2$ -graded associative algebra over $\mathbb{C}$ with generators $t_{ij}^{(r)}$ and certain relations described below. We define the formal power series

$$t_{ij}(u) = \delta_{ij} + t_{ij}^{(1)}u^{-1} + t_{ij}^{(2)}u^{-2} + \dots,$$

and a matrix

$$T(u) = \sum_{i,j=1}^{m+n} t_{ij}(u) \otimes E_{ij} (-1)^{\bar{j}(\bar{i}+1)}, \quad (1)$$

where E_{ij} is the standard elementary matrix and $\bar{i}$ is the parity of the index i . In analogy with the usual Yangian $Y(\mathfrak{gl}_n)$ (see for example [2], [5], [6]), the defining relations are then expressed by the matrix product

$$R(u-v)T_1(u)T_2(v) = T_2(v)T_1(u)R(u-v)$$

where

$$R(u-v) = 1 - \frac{1}{(u-v)}P_{12}$$

and P_{12} is the permutation matrix:

$$P_{12} = \sum_{i,j=1}^{m+n} E_{ij} \otimes E_{ji} (-1)^{\bar{j}}.$$

Then we have the following equivalent form of the defining relations:

$$[t_{ij}(u), t_{kl}(v)] = \frac{(-1)^{\bar{i}\bar{j} + \bar{i}\bar{k} + \bar{j}\bar{k}}}{(u-v)} (t_{kj}(u)t_{il}(v) - t_{kj}(v)t_{il}(u)).$$

*lucyg@maths.usyd.edu.au

¹The author would like to acknowledge the guidance and support of her PhD supervisor Alexander Molev.

Throughout this article we will observe the following notation for entries of the inverse matrix of $T(u)$:

$$T(u)^{-1} =: (t'_{ij}(u))_{i,j=1}^n.$$

A straightforward calculation yields the following relation in $Y(\mathfrak{gl}_{m|n})$:

$$[t_{ij}(u), t'_{kl}(v)] = \frac{(-1)^{\overline{i}\overline{j} + \overline{i}\overline{k} + \overline{j}\overline{k}}}{(u-v)} \cdot (\delta_{kj} \sum_{s=1}^{m+n} t_{is}(u) t'_{sl}(v) - \delta_{il} \sum_{s=1}^{m+n} t'_{ks}(v) t_{sj}(u)). \quad (2)$$

2 Gauss Decomposition of $T(u)$

In [1], the Drinfeld presentation is described in terms of the quasideterminants of Gelfand and Retakh ([3], [4]). In this article we make use of the analogous set of generators of the Yangian $Y(\mathfrak{gl}_{m|n})$. First we recall the definition of the quasideterminants and some conventional notation.

Definition 2.1. *Let X be a square matrix over a ring with identity such that its inverse matrix X^{-1} exists, and such that its j th entry is an invertible element of the ring. Then the ij th quasideterminant of X is defined by the formula*

$$|X|_{ij} = ((X^{-1})_{ji})^{-1}.$$

It is sometimes convenient to adopt the following alternative notation for the quasideterminants:

$$|X|_{ij} =: \begin{vmatrix} x_{11} & \cdots & x_{1j} & \cdots & x_{1n} \\ & \ddots & & \ddots & \\ x_{i1} & \cdots & \boxed{x_{ij}} & \cdots & x_{in} \\ & \ddots & & \ddots & \\ x_{n1} & \cdots & x_{nj} & \cdots & x_{nn} \end{vmatrix}.$$

The matrix $T(u)$ defined in (1) has the following Gauss decomposition in terms of quasideterminants (by Theorem 4.96 in [3]; see §5 in [1]):

$$T(u) = F(u)D(u)E(u)$$

for unique matrices

$$\begin{aligned} D(u) &= \begin{pmatrix} d_1(u) & & \cdots & 0 \\ & d_2(u) & & \vdots \\ \vdots & & \ddots & \\ 0 & \cdots & & d_{m+n}(u) \end{pmatrix}, \\ E(u) &= \begin{pmatrix} 1 & e_{12}(u) & \cdots & e_{1,m+n}(u) \\ & \ddots & & e_{2,m+n}(u) \\ & & \ddots & \vdots \\ 0 & & & 1 \end{pmatrix}, \\ F(u) &= \begin{pmatrix} 1 & & \cdots & 0 \\ f_{21}(u) & \ddots & & \vdots \\ \vdots & & \ddots & \\ f_{m+n,1}(u) & f_{m+n,2}(u) & \cdots & 1 \end{pmatrix}, \end{aligned}$$

where

$$\begin{aligned}
d_i(u) &= \begin{vmatrix} t_{11}(u) & \cdots & t_{1,i-1}(u) & t_{1i}(u) \\ \vdots & \ddots & & \vdots \\ t_{i1}(u) & \cdots & t_{i,i-1}(u) & \boxed{t_{ii}(u)} \end{vmatrix}, \\
e_{ij}(u) &= d_i(u)^{-1} \begin{vmatrix} t_{11}(u) & \cdots & t_{1,i-1}(u) & t_{1j}(u) \\ \vdots & \ddots & \vdots & \vdots \\ t_{i-1,i}(u) & \cdots & t_{i-1,i-1}(u) & t_{i-1,j}(u) \\ t_{i1}(u) & \cdots & t_{i,i-1}(u) & \boxed{t_{ij}(u)} \end{vmatrix}, \\
f_{ji}(u) &= \begin{vmatrix} t_{11}(u) & \cdots & t_{1,i-1}(u) & t_{1i}(u) \\ \vdots & \ddots & \vdots & \vdots \\ t_{i-1,1}(u) & \cdots & t_{i-1,i-1}(u) & t_{i-1,i}(u) \\ t_{ji}(u) & \cdots & t_{j,i-1}(u) & \boxed{t_{ji}(u)} \end{vmatrix} d_i(u)^{-1}.
\end{aligned}$$

It is easy to recover each generating series $t_{ij}(u)$ by multiplying together and taking commutators of $d_i(u)$; $1 \leq i \leq m+n$, and $e_i(u) := e_{i,i+1}(u)$, $f_i(u) = f_{i+1,i}(u)$; $1 \leq i < m+n$ (see §5 of [1]). Thus the Yangian $Y(\mathfrak{gl}_{m|n})$ is generated by the coefficients of the latter.

2.1 Some Useful Maps

Here we define some automorphisms of the Yangian $Y(\mathfrak{gl}_{m|n})$ and homomorphisms between Yangians, so that we may refer to them in the next section.

Let $\omega_{m|n} : Y(\mathfrak{gl}_{m|n}) \rightarrow Y(\mathfrak{gl}_{m|n})$ be the automorphism defined by

$$\omega : T(u) \mapsto T(-u)^{-1}.$$

Let $\tau : Y(\mathfrak{gl}_{m|n}) \rightarrow Y(\mathfrak{gl}_{m|n})$ be the anti-automorphism defined by

$$\tau(t_{ij}(u)) = t_{ji}(u) \times (-1)^{\bar{i}(\bar{j}+1)}.$$

Let $\rho_{m|n} : Y(\mathfrak{gl}_{m|n}) \rightarrow Y(\mathfrak{gl}_{n|m})$ be the isomorphism defined by

$$\rho_{m|n}(t_{ij}(u)) = t_{m+n+1-i, m+n+1-j}(-u).$$

Let $\varphi_{m|n} : Y(\mathfrak{gl}_{m|n}) \hookrightarrow Y(\mathfrak{gl}_{m+k|n})$ be the inclusion which sends each generator $t_{ij}^{(r)} \in Y(\mathfrak{gl}_{m|n})$ to the generator $t_{k+i, k+j}^{(r)}$ in $Y(\mathfrak{gl}_{m+k|n})$.

Finally, let $\psi_k : Y(\mathfrak{gl}_{m|n}) \rightarrow Y(\mathfrak{gl}_{m+k|n})$ be the injective homomorphism defined by

$$\psi_k = \omega_{m+k|n} \circ \varphi_{m|n} \circ \omega_{m|n}. \quad (3)$$

This last homomorphism is useful for studying quasideterminants so we discuss it in some detail with the following remarks.

Remark 2.1. We can calculate $\psi_k(t_{ij}(u))$ explicitly for any $1 \leq i, j \leq m+n$ (see Lemma 4.2 of [1]) :

$$\psi_k(t_{ij}(u)) = \begin{vmatrix} t_{11}(u) & \cdots & t_{1k}(u) & t_{1,k+j}(u) \\ \vdots & \ddots & \vdots & \vdots \\ t_{k1}(u) & \cdots & t_{kk}(u) & t_{k,k+j}(u) \\ t_{k+i,1}(u) & \cdots & t_{k+i,k}(u) & \boxed{t_{k+i,k+j}(u)} \end{vmatrix}.$$

In particular, this means that for $k \geq 1$, we have $\psi_k(d_1(u)) = d_{k+1}(u)$, $\psi_k(e_1(u)) = e_{k+1}(u)$, and $\psi_k(f_1(u)) = f_{k+1}(u)$.

Furthermore, by (3), we have for any $k, l \geq 1$ that $\psi_k \circ \psi_l = \psi_{k+l}$, so we may generalise this observation to give for instance $\psi_k(d_l(u)) = d_{k+l}(u)$.

Remark 2.2. Notice that the map ψ_k sends $t'_{ij}{}^{(r)} \in Y(\mathfrak{gl}_{m|n})$ to the element $t'_{k+i,k+j}{}^{(r)}$ in $Y(\mathfrak{gl}_{m+k|n})$. Thus the subalgebra $\psi_k(Y(\mathfrak{gl}_{m|n}))$ is generated by the elements $\{t'_{k+s,k+t}{}^{(r)}\}_{s,t=1}^n$. Then, by (2), all elements of this subalgebra commute with those of the subalgebra generated by the elements $\{t'_{ij}{}^{(r)}\}_{i,j=1}^k$.

By Remark 2.1, this implies in particular that for any $i, j \geq 1$, the quasideterminants $d_i(u)$ and $d_j(v)$ commute.

3 The Quantum Berezinian

The quantum Berezinian was defined by Nazarov [7] and plays a similar role in the study of the Yangian $Y(\mathfrak{gl}_{m|n})$ as the quantum determinant does in the case of the Yangian $Y(\mathfrak{gl}_n)$ (see [5]).

Definition 3.1. The quantum Berezinian is the following power series with coefficients in the Yangian $Y(\mathfrak{gl}_{m|n})$:

$$\begin{aligned} b_{m|n}(u) &:= \sum_{\tau \in S_m} \text{sgn}(\tau) t_{\tau(1)1}(u) t_{\tau(2)2}(u-1) \cdots t_{\tau(m)m}(u-m+1) \\ &\quad \times \sum_{\sigma \in S_n} \text{sgn}(\sigma) t'_{m+1,m+\sigma(1)}(u-m+1) \cdots t'_{m+n,m+\sigma(n)}(u-m+n) \end{aligned}$$

The first part of this expression for $b_{m|n}(u)$ is quite special and so is given its own notation:

$$C_m(u) := \sum_{\tau \in S_m} \text{sgn}(\tau) t_{\tau(1)1}(u) t_{\tau(2)2}(u-1) \cdots t_{\tau(m)m}(u-m+1).$$

It is clear that $C_m(u)$ is an element of the subalgebra of $Y(\mathfrak{gl}_{m|n})$ generated by the set $\{t'_{ij}{}^{(r)}\}_{1 \leq i,j \leq m; r \geq 0}$. This subalgebra is isomorphic to the Yangian $Y(\mathfrak{gl}_m)$ associated to the Lie algebra $\mathfrak{gl}_m$ by the inclusion $Y(\mathfrak{gl}_m) \rightarrow Y(\mathfrak{gl}_{m|n})$ which send each generator $t'_{ij}{}^{(r)}$ in $Y(\mathfrak{gl}_m)$ to the generator of the same name in $Y(\mathfrak{gl}_{m|n})$. Moreover, $C_m(u)$ is in fact the image under this map of the *quantum determinant* of the smaller Yangian $Y(\mathfrak{gl}_m)$ (see [1], [5]). Then it is well known (see Theorem 2.32 in [6]) that we have the alternative expression:

$$C_m(u) = d_1(u) d_2(u-1) \cdots d_m(u-m+1).$$

We can extend this observation as follows:

Theorem 1. We have the following alternative expression for the quantum Berezinian:

$$\begin{aligned} b_{m|n}(u) &= d_1(u) d_2(u-1) \cdots d_m(u-m+1) \\ &\quad \times d_{m+1}(u-m+1)^{-1} \cdots d_{m+n}(u-m+n)^{-1}. \end{aligned}$$

Proof. Notice that the second part of the expression for $b_{m|n}(u)$ in Definition 3.1 is the image under the isomorphism $\rho_{n|m} \circ \omega_{n|m} : Y(\mathfrak{gl}_{n|m}) \rightarrow Y(\mathfrak{gl}_{m|n})$ of

$$\sum_{\sigma \in S_n} \text{sgn}(\sigma) t_{n,\sigma(n)}(u-m+1) \cdots t_{2,\sigma(2)}(u-m+n-1) t_{1,\sigma(1)}(u-m+n) \quad (4)$$

where in this expression (4) we are following the usual convention for denoting generators in the Yangian $Y(\mathfrak{gl}_{n|m})$. We recognise (by comparing with (8.3) of [1] for example) that the expression (4) is in fact $C_n(u-m+n)$, the image of the quantum determinant of $Y(\mathfrak{gl}_n)$ under the natural inclusion $Y(\mathfrak{gl}_n) \hookrightarrow Y(\mathfrak{gl}_{n|m})$. So in order to verify the claim we must calculate the image of $C_n(u-m+n)$ under this map explicitly in terms of our quasideterminants $d_i(v)$. Applying Proposition 1.6 of [4], we find that the image of $d_i(v)$ in $Y(\mathfrak{gl}_{n|m})$ is $(d_{m+n+1-i}(v))^{-1}$ in $Y(\mathfrak{gl}_{m|n})$. This gives the desired result. $\square$

The following theorem is a result of Nazarov [7]. We give a new proof.

Theorem 2. *The coefficients of the quantum Berezinian (3.1) are central in the algebra $Y(\mathfrak{gl}_{m|n})$.*

Proof. By Remark 2.2, we already know that the quantum Berezinian $b_{m|n}(u)$ commutes with $d_i(v)$ for $1 \leq i \leq m+n$. So our problem reduces to showing that $b_{m|n}(u)$ commutes with $e_i(v)$ and with $f_i(v)$ for each i between 1 and $m+n-1$. We proceed by breaking this problem into three cases.

Case 1: $1 \leq i \leq m-1$. In this case, $e_i(v)$ commutes with $C_m(u) = d_1(u) \cdots d_m(u-m+1)$ by Theorem 7.2 in [1]. On the other hand, $e_i(v)$ is an element of the subalgebra generated by $\{t_{jk}^{(r)}\}_{1 \leq j,k \leq m}$ and thus by Remark 2.2 commutes with $d_{m+s}(u-m+s)^{-1} = t'_{m+s,m+s}(u-m+s)$ for $1 \leq s \leq n$. Now we may use the anti-automorphism τ to show that $f_i(v)$ also commutes with the quantum Berezinian in this case because $\tau(e_i(v)) = f_i(v)$ for $1 \leq i \leq m-1$.

Case 2: $m+1 \leq i \leq m+n-1$. Applying Propositions 1.6 and 1.4 of [4] in turn to $f_i(v)$, we find an alternative expression:

$$f_i(v) = - \left| \begin{array}{cccc} \boxed{t'_{i+1,i}(v)} & t'_{i+1,i+2}(v) & \cdots & t'_{i+1,m+n}(v) \\ t'_{i+2,i}(v) & t'_{i+2,i+2}(v) & \cdots & \\ \vdots & \vdots & \ddots & \vdots \\ t'_{m+n,i}(v) & t'_{m+n,i+2}(v) & \cdots & t'_{m+n,m+n}(v) \end{array} \right| \left| \begin{array}{ccc} \boxed{t'_{i+1,i+1}(v)} & \cdots & t'_{i+1,m+n}(v) \\ \vdots & & \vdots \\ t'_{m+n,i+1}(v) & \cdots & t'_{m+n,m+n}(v) \end{array} \right|^{-1}.$$

Thus, we find that for $m+1 \leq i \leq m+n-1$,

$$\rho_{n|m} \circ \omega_{n|m}(-f_{m+n-i}(v)) = e_i(v),$$

and similarly

$$\rho_{n|m} \circ \omega_{n|m}(-e_{m+n-i}(v)) = f_i(v).$$

We apply this isomorphism to the results of Case 1 in the Yangian $Y(\mathfrak{gl}_{n|m})$. This shows that $e_i(v)$ and $f_i(v)$ commute with the quantum Berezinian in the case where $m+1 \leq i \leq m+n-1$.

Case 3: $i=m$. We begin by considering the Yangian $Y(\mathfrak{gl}_{1|1})$. For this algebra we have that the quantum Berezinian is $b_{1|1}(u) = d_1(u)d_2(u)^{-1}$ and we would like to show that it commutes with $e_1(v)$ and $f_1(v)$. So it will suffice to show

$$d_1(u)e_1(v)d_2(u) = d_2(u)e_1(v)d_1(u). \quad (5)$$

We have

$$\begin{pmatrix} t_{11}(u) & t_{12}(u) \\ t_{21}(u) & t_{22}(u) \end{pmatrix} = \begin{pmatrix} d_1(u) & d_1(u)e_1(u) \\ f_1(u)d_1(u) & f_1(u)d_1(u)e_1(u) + d_2(u) \end{pmatrix} \quad (6)$$

$$\begin{pmatrix} t'_{11}(v) & t'_{12}(v) \\ t'_{21}(v) & t'_{22}(v) \end{pmatrix} = \begin{pmatrix} d_1(v)^{-1} + e_1(v)d_2(v)^{-1}f_1(v) & -e_1(v)d_2(v)^{-1} \\ -d_2(v)^{-1}f_1(v) & d_2(v)^{-1} \end{pmatrix}. \quad (7)$$

An application of (2) gives

$$(u-v)[t_{11}(u), t'_{12}(v)] = t_{11}(u)t'_{12}(v) + t_{12}(u)t'_{22}(v).$$

Substituting in the expressions from (6) and (7) then cancelling $d_2(v)$, this gives

$$(u-v)[d_1(u), e_1(v)] = d_1(u)(e_1(v) - e_1(u)).$$

Similarly, by considering the commutator $[t_{12}(u), t'_{22}(v)]$, we derive the relation

$$(u-v)[d_2(u), e_1(v)] = d_2(u)(e_1(v) - e_1(u)).$$

We rewrite these relations to find

$$\begin{aligned} (u-v)e_1(v)d_1(u) &= (u-v-1)d_1(u)e_1(v) + d_1(u)e_1(u), \\ (u-v)e_1(v)d_2(u) &= (u-v-1)d_2(u)e_1(v) + d_2(u)e_1(u), \end{aligned}$$

and by considering these expressions we see that (5) holds. Similarly, the quantum Berezinian commutes with $f_1(v)$.

Now we return our attention to the general Yangian $Y(\mathfrak{gl}_{m|n})$. By similar appeals to Remark 2.2 as in the first case, we see that $e_m(v)$ and $f_m(v)$ commute with $d_1(u) \cdots d_{m-1}(u - m + 2)$ and with $d_{m+2}(u - m + 2)^{-1} \cdots d_{m+n}(u - m + n)^{-1}$. So we need only show they commute with $d_m(u - m + 1)d_{m+1}(u - m + 1)^{-1}$. This follows immediately when we apply the homomorphism ψ_{m-1} to the identity (5) in $Y(\mathfrak{gl}_{1|1})$. $\square$

References

- [1] Jonathan Brundan and Alexander Kleshchev. Parabolic presentations of the Yangian $Y(\mathfrak{gl}_n)$. *To appear in Commun. Math. Phys.*, 2004. Preprint: math.QA/0407011.
- [2] Vyjayanthi Chari and Andrew Pressley. *A guide to quantum groups*. Cambridge University Press, Cambridge, 1994.
- [3] I. Gelfand, S. Gelfand, V. Retakh, and R.L. Wilson. Quasideterminants. *to appear in Advances in Mathematics*, 2004. Preprint: arXiv:math.QA/0208146.
- [4] I. M. Gelfand and V. S. Retakh. Determinants of matrices over noncommutative rings. *Funct. Anal. Appl.*, 25(2):91–102, 1991.
- [5] A. Molev, M. Nazarov, and G. Olshanskii. Yangians and classical Lie algebras. *Russian Math. Surveys*, 51:205–282, 1996.
- [6] A. I. Molev. Yangians and their applications. In *Handbook of algebra, Vol. 3*, pages 907–959. North-Holland, Amsterdam, 2003.
- [7] M. L. Nazarov. Quantum Berezinian and the classical Capelli identity. *Lett. Math. Phys.*, 21(2):123–131, 1991.